

Maquoketa Community School District



Kristy Haxmeier
Business Manager

612 South Vermont Street
Maquoketa, Iowa 52060-3460
Phone (563) 652-4984
Website: maquoketaschools.org

Mike Wing
Director of Transportation &
Manager of Support Operations

Jane Schmidt
Director of Professional Growth
& Student Learning

Mr. Christopher Hoover
Superintendent of Schools

Dustin Cogan
Technology Director

January 25, 2018

Dear District and Community Patrons,

There have been 2 suspect, unconfirmed cases of mumps in the Maquoketa School system reported this week. When we receive more information about either of these cases we will let you know. Please see below what the symptoms of the mumps are and what you should do if you suspect that you, or your child, might have the disease.

What are the symptoms of mumps?

The most common symptoms of mumps include:

- Fever
- Headache
- Muscle aches
- Tiredness
- Loss of appetite
- Swollen and tender salivary glands under the ears on one or both sides (parotitis)

Some people, however, have very mild or no symptoms and often they do not know they have the disease.

How is mumps spread?

Mumps is a contagious disease caused by a virus. It spreads through saliva or mucus from the mouth, nose, or throat.

An infected person can spread the virus by:

- kissing, coughing, sneezing, or talking
- sharing items, such as cups or eating utensils, with others
- touching objects or surfaces with unwashed hands that are then touched by others

Mumps likely spreads before the salivary glands begin to swell and up to five days after the swelling begins.

What should you do if you think you may have mumps?

Contact your healthcare provider who may wish to test for mumps.

What should people infected with mumps do to prevent spreading it to others?

When you have mumps, you should avoid prolonged, close contact with other people until five days after symptoms began or until you are well, whichever is longer. You should stay home and limit

OUR MISSION: To ensure all students and staff are learning at high levels.

OUR VISION: To develop a culture of collaboration and communication that supports continuous improvement for all.

contact with the people you live with; for example, sleep by yourself in a separate room if you can. Staying home while sick with mumps is an important way to avoid spreading the virus to other people.

Can people who have been vaccinated with MMR still be infected with mumps?

MMR vaccine prevents most, but not all, cases of mumps and complications caused by the disease. Two doses of the vaccine are about 88 percent effective at preventing mumps; one dose is about 78 percent effective.

People who have received two doses of the MMR vaccine are about nine times less likely to get mumps than unvaccinated people who have the same exposure to mumps virus. However, some people who receive two doses of MMR can still get mumps, especially if they have prolonged, close contact with someone who has the disease. If a vaccinated person does get mumps, they will likely have less severe illness than an unvaccinated person.

Should everyone get a third dose of MMR vaccine to protect against mumps?

When mumps cases are occurring at schools, colleges or universities, national criteria are used to determine when a third dose of MMR should be recommended. No other colleges or universities in Iowa have met the national criteria; therefore, a third dose of MMR vaccine has not been recommended by public health officials.

What should schools do if mumps cases have been identified in their facilities?

Schools should reinforce that persons infected with mumps should not go back to school, work, or other public places until five days after symptoms began or until they are well, whichever is longer.

What should schools do if students or staff are exposed to a person with mumps?

IDPH does not recommend excluding asymptomatic exposed persons from school or work. Persons who may have been exposed should be educated on the signs and symptoms of mumps disease, and should seek medical attention as soon as any of these symptoms begin.

What should every school do to decrease or help prevent mumps transmission?

Schools should promote healthy habits among students and staff, such as:

- Covering your mouth and nose with a tissue when you cough or sneeze, and putting your used tissue in the trash can. If you don't have a tissue, cough or sneeze into your upper sleeve or elbow, not your hands.
- Washing your hands often with soap and water.
- Avoiding sharing drinks or eating utensils.
- Disinfecting frequently touched surfaces, such as doorknobs, tables, counters.
- Be up-to-date on the MMR vaccine.

Should schools perform extra cleaning if a mumps infected person is identified in their facility?

Routine cleaning procedures are sufficient. Special attention should be paid to commonly touched surfaces.

OUR MISSION: To ensure all students and staff are learning at high levels.

OUR VISION: To develop a culture of collaboration and communication that supports continuous improvement for all.